

## PART III: CONSUMER INFORMATION

### Pr **PROPAFENONE** propafenone hydrochloride, film-coated tablets

This leaflet is PART III of a three-part "Product Monograph" published when PROPAFENONE was approved for sale in Canada and is designed specifically for consumers. This leaflet is a summary and will not tell you everything about PROPAFENONE. Contact your doctor or pharmacist if you have any questions about the drug.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

- PROPAFENONE is used to control certain types of irregular heartbeats (arrhythmias).

##### What it does:

PROPAFENONE is a heart rate regulating agent. It acts on the metabolism of the heart muscles to block some of the irregular heartbeats. It also acts as a local anaesthetic, blocks the sodium current and slows down the potential of heart muscles reacting fast.

##### When it should not be used:

PROPAFENONE should not be used if:

- you are allergic to any component of PROPAFENONE, including active ingredients and non-active ingredients;
- you have certain serious heart conditions or incidence of heart attack within the last 3 months;
- you have serious liver failure;
- you have certain respiratory conditions;
- you have myasthenia gravis.
- you are younger than 18

##### What the medicinal ingredient is:

propafenone hydrochloride

##### What the important non-medicinal ingredients are:

croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, maize starch, microcrystalline cellulose, polyethylene glycol 400 and 6000, purified water, titanium dioxide

*For a full listing of non-medicinal ingredients see PART I of the Product Monograph.*

##### What dosage forms it comes in:

PROPAFENONE is available as film-coated tablets in the following strengths: 150 mg and 300 mg.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

- PROPAFENONE is intended for use only in patients with life-threatening irregular heartbeats (arrhythmias). Most anti-arrhythmic drugs have the potential to cause dangerous arrhythmias; some have been shown to be associated with an increase of sudden death. Your doctor will tell you about the risk and benefits of anti-arrhythmic therapy.

#### BEFORE you use PROPAFENONE talk to your doctor or pharmacist if:

- you have a family history of sudden cardiac death or suffer from Brugada Syndrome;
- you are pregnant or planning to become pregnant, or you are breast-feeding;
- you have any heart disease;
- you have abnormal blood cell counts;
- you have abnormal liver function;
- you have neuromuscular disease (e.g. myasthenia gravis);
- you have kidney disease;
- you have allergies to this drug or any of its ingredients.
- you perform tasks which require special attention (for example, driving automobile or operating dangerous machinery) because blurred vision, dizziness, fatigue and low blood pressure are common side effects associated with the administration of PROPAFENONE.

## INTERACTIONS WITH THIS MEDICATION

#### Drugs that may interact with PROPAFENONE include:

- beta-blockers (e.g. propranolol, metoprolol);
- digoxin, venlafaxine, rifampin, cimetidine, quinidine, ketoconazole, erythromycin, amiodarone, phenobarbital;
- anticoagulants (e.g. warfarin);
- certain local anesthetics (e.g. lidocaine);
- certain antidepressants of the tricyclic group (e.g. desipramine), and other antidepressants (e.g. fluoxetine, paroxetine, fluvoxamine);
- some medication that can affect your immune system (e.g. cyclosporine);
- some HIV-antiviral medication (e.g. ritonavir, lopinavir/ritonavir);
- grapefruit juice.

## PROPER USE OF THIS MEDICATION

#### Usual dose:

Dosage must be individualized. The usual adult dose of PROPAFENONE is 150 mg which is to be taken every 8 hours, however your doctor may decide on different individual dosing.

The film-coated tablets should be swallowed whole (without chewing) with liquid. Recommended to be taken with food.

**Overdose:**

If you or someone you know accidentally takes more than stated dose, contact your doctor or Regional Poison Control Centre immediately or go to the nearest hospital with the tablets. Tell your doctor or hospital how much was taken. Treat even small overdoses seriously.

**Missed Dose:**

If you forget to take one tablet, take another as soon as you remember, unless it is almost time for your next dose. If it is, do not take the missed tablet at all.

Never double-up on a missed dose.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Along with its needed effects, a medicine may cause some unwanted effects. These are referred to as “side effects”. Although not all of these side effects may occur, if they do occur they may need medical attention.

The most common side effects with PROPAFENONE are dizziness, feeling sick (nausea), vomiting, unusual taste, constipation. Other less common side effects may include headaches, blurred vision, difficulty in sleeping, tremor, drowsiness, dyspepsia, dry mouth, loss of appetite, abdominal pain/cramping, flatulence, tiredness, skin rash, weakness, chest pain, anxiety, severe sweating and pain in the joints.

**Check with your physician or pharmacist if you experience any unexpected effects, or are concerned by the above side effects.**

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom/effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist as soon as possible	
	Only if severe	In all cases		
Very common	Fast heartbeat, irregular heartbeats		√	
Common	chest pain,		√	
	dizziness, lightheadedness, fainting		√	
	liver problems (e.g., yellowing skin or eyes, prolonged vomiting and nausea or abdominal pain)		√	
	bleeding problem (excessive bruising, easy bleeding)		√	
Not common	Abnormal muscular control (ataxia)		Ⓣ	
Unknown	Heart rate reduced		Ⓣ	
	Convulsion, movement disorders (extrapyramidal symptoms), restlessness		Ⓣ	

**This is not a complete list of side effects. For any unexpected effects while taking PROPAFENONE, contact your doctor or pharmacist.**

Check with your pharmacist or doctor **immediately**, if you experience any of the above symptoms of the serious side effects.

**HOW TO STORE IT**

Keep PROPAFENONE and all other medicines out of reach of children.

PROPAFENONE tablets should be stored at 15° to 25°C, protected from light and moisture.

Do not take your tablets after the expiry date shown on the label.

It is important to keep the PROPAFENONE tablets in the original package.

**Reporting Side Effects**

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

**3 ways to report:**

- Online at [MedEffect](#);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:

- Fax to 1-866-678-6789 (toll-free), or
- Mail to: Canada Vigilance Program  
Health Canada, Postal Locator  
0701E  
Ottawa, ON  
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at [MedEffect](#).

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice. Program does not provide medical advice.*

**MORE INFORMATION**

The full Product Monograph, prepared for health professionals can be found by contacting the sponsor, Sanis Health Inc., at: 1-866-236-4076.

This leaflet was prepared by Sanis Health Inc., 1 President's Choice Circle, Brampton, Ontario, L6Y 5S5.

Prepared on April 18, 2017